



hCAR·CD19-T Cell		
一、产品信息		
产品编号	产品名称	产品规格
VGT-930-26A2	hCAR-CD19-T 细胞	1 × 10 ⁷ 细胞
二、产品介绍		
Generation	Second	
Receptor Construction	Anti CD19 scFv (FMC63)-hCD28-hCD3ζ	
Marker	CAR	
Vector Type/Length	Retrovirus / 9 kb	
Description of Signaling Cassettes	<p>CD28:</p> <p>CD28 (Cluster of Differentiation 28) is one of the proteins expressed on T cells that provide co-stimulatory signals required for T cell activation and survival. CD28 is the receptor for CD80 (B7.1) and CD86 (B7.2) proteins which are expressed on antigen-presenting cells (APC). CD28 modulates the primary TCR/CD3ζ signal in a different fashion than the late costimulatory elements OX40 and 4-1BB. CD28 enhances the expression of downstream regulators that impact on T-cell proliferation, death, differentiation, and effector functions. CAR+ T cells containing the CD28 endodomain showed strikingly enhanced sustained T cell activation, growth, survival. And CD28 results in a brightly expressed, stable receptor as the transmembrane domain. Including CD28 costimulatory domains in CARs led to enhanced anti-malignancy efficacy.</p>	





维根生物科技有限公司

Vigen Biotechnology (Zhenjiang) Co., Ltd

VigenCAR®

Tel: 183 628 99236

E-mail: 253540644@qq.com

	<p>CD3ζ:</p> <p>CD3ζ, also known as T-cell receptor zeta, which together with T-cell receptor and CD3γ, δ, ε chain, forms the TCR-CD3 complex. ζ was expressed independently from the complex. The zeta chain plays an important role in coupling antigen recognition to several intracellular signal-transduction pathways. CD3-zeta, which contains 3 ITAMs, is the most commonly used endodomain component of CARs. It transmits an activation signal to the T cell after antigen is bound. CD3-zeta may not provide a fully competent activation signal and additional co-stimulatory signaling is needed. For example, chimeric CD28 and OX40 can be used with CD3-zeta to transmit a proliferative/survival signal, or all three can be used together.</p>	
Effector cell type	T cell (human)	
Target	CD19	
Synonyms	B4; CVID3	
Target Gene ID	930	
Targeting Diseases	Chronic lymphocytic leukemia	
三、产品组分		
组分编号	组分名称	产品规格
VGT-930-26A2	hCAR-CD19-T 细胞	1 × 10 ⁷ 细胞
VGT-0000-18A4	ATC 细胞（未转导活化 T 细胞）	1 × 10 ⁷ 细胞
VGA-0013-0100	人 T 细胞完全培养基	100 mL
VGA-0036-0200	重组人 IL-2 溶液（1000x）	100 μL



www.vigenbio.com



Technical Support



维根生物科技有限公司
Vigen Biotechnology (Zhenjiang) Co., Ltd

VigenCAR®

Tel: 183 628 99236
E-mail: 253540644@qq.com

运输与保存:

干冰运输。收到后如不立即复苏，请立即转入-80°C 保存或液氮储存。

四、限制性使用许可

- 1) 本产品仅供购买方单方使用，不得向任何第三方赠送、转让或销售；
- 2) 本产品仅供实验室和体外研究使用，不能用于任何临床检验、诊断、治疗，或任何非法用途；
- 3) 收货 48 h 内如若发现异常，请及时联系售后，逾期视为收货良好；
- 4) 请严格按照本说明书操作，否则造成细胞失活等情形，不予提供补发服务。



www.vigenbio.com



Technical Support